

A New Photometric Model of the Galactic Bar using Red Clump Giants

Liang Cao¹, Shude Mao^{1,2}, David Nataf³, Nicholas J. Rattenbury², Andrew Gould³

¹ *National Astronomical Observatories, 20A Datun Road, Chinese Academy of Sciences, Beijing, 100012, China*

² *Jodrell Bank Centre for Astrophysics, The University of Manchester, Alan Turing Building, Manchester M13 9PL, UK*

³ *Department of Astronomy, The Ohio State University, 140 West 18th Ave., Columbus, OH 43210, USA*

Accepted Received ; in original form.....

ABSTRACT

We present a study of the luminosity density distribution of the Galactic bar using number counts of red clump giants (RCGs) from the OGLE-III survey. The data were recently released by Nataf et al. (2012) for 9019 fields towards the bulge and have 2.94×10^6 RC stars over a viewing area of 90.25 deg^2 . The data include the number counts, mean distance modulus (μ), dispersion in μ and full error matrix. We fit the data with several tri-axial parametric models, after excluding the bad sightlines, 2.65×10^6 RC stars are remaining with the sky coverage of 81.47 deg^2 . We use the Markov Chain Monte Carlo (MCMC) method to explore the parameter space and find that the best-fit model is close to exponential (E_2 model), with the distance to the GC is 8.17 kpc, the ratio of semi-major and semi-minor bar axis scale lengths in the Galactic plane x_0, y_0 , and vertical bar scale length z_0 , is $x_0 : y_0 : z_0 \approx 1.00 : 0.41 : 0.38$ (close to being prolate). The bar major axis is ~ 0.68 kpc and has an angle of between 29° and 32° , slightly larger than the value obtained from a similar study based on OGLE-II data. The number of estimated RC stars is 2.51×10^6 , which is systematically lower than the observed value. We subtract the smooth parametric model from the observed counts and find that the residuals are consistent with the presence of an X-shaped structure in the Galactic centre, the excess to the estimated mass content is $\sim 5.3\%$. The results can be used as a key ingredient to construct new density models of the Milky Way and will have implications on the predictions of the optical depth to gravitational microlensing and the patterns of hydrodynamical gas flow in the Milky Way.

Key words: Galaxy - structure: Galaxy - stellar content - Galaxy: bulge - Galaxy: centre

1 INTRODUCTION

Most spiral galaxies in the local Universe are barred. Galactic bars are often straight and their presence appears to be intimately related to star formation. Rings are often seen in these galaxies and they are likely related to resonances in a rotating barred potential. Thus barred galaxies offer exciting opportunities to understand gas and dynamical processes in spiral galaxies (see e.g. Sellwood & Wilkinson 1993; Athanassoula 2012 for reviews). It is now generally accepted that the inner region of our own Galaxy also hosts a tri-axial, bar-like structure. Observational evidence for a bar has arisen from several sources, such as the studies of gas kinematics (e.g. Binney et al. 1991), surface brightness (e.g. Blitz & Spergel 1991), star counts (e.g. Nakada et al. 1991; Stanek et al. 1994) and microlensing (e.g. Udalski et al. 1994; see Gerhard 2002 for a review).

Various tracers have been used to constrain stellar density models of the inner Galaxy. Dwek et al. (1995) used the COBE-DIRBE multi-wavelength observations of the Galactic centre (Weiland et al. 1994) to constrain several multi-parameter analytic bar models. Stanek et al. (1997) first used red clump giants (RCGs) to constrain bar models (see also Babusiaux & Gilmore 2005 and Nishiyama et al. 2005 who traced the bulge RCG population in the infrared). Rattenbury et al. (2007) applied the method of Stanek et al. (1997) to OGLE-II data and updated the bar parameters using the RCG population.

Unfortunately these studies leave many bar parameters uncertain. This may reflect the intrinsic complexities of the Galactic bar. For example, there has been a fair amount of discussion about the existence of a nuclear bar. Nishiyama et al. (2005) found a clear change of slope in the red clump giants around $|l| = 4^\circ$, which was confirmed recently by the VVV survey (Gonzalez et al. 2011). This was interpreted as the existence of a nuclear bar. However, numerically simulated bars appear to be able to reproduce the changes in the slope without any nuclear structure (Gerhard & Martinez-Valpuesta 2012). With more and more dynamical data emerging from several kinematic surveys (e.g., BRAVA, Howard et al. 2008, ARGOS, Freeman et al. 2013, and APOGEE, Nidever et al. 2012), it appears increasingly important to construct a more accurate photometric model, which will serve as a key ingredient in self-consistent dynamical modelling (Wang et al. 2012; Long et al. 2013).

The outline of this paper is as follows. In §2 we summarize the observational data used, as released by Nataf et al. (2012); in §3 we describe the methodology used in this analysis; in §4, we apply our methods to the OGLE-III data, and present the main results of this work. In §5 we discuss our results and compare them with earlier work.

2 DATA SAMPLE

In this paper, we explore the OGLE-III observation data recently released by Nataf et al. (2012) who carefully studied the extinction and the red clump giant population using both OGLE and 2MASS data. The RC is a prominent, well-populated, and localized feature of Galactic bulge CMDs (Terndrup et al. 1998; Stanek et al. 1994). RCGs are very good standard candles with $M_I = -0.12$ and an intrinsic dispersion of 0.09 mag in luminosity at the estimated age and metallicity distribution of the bulge stellar population (Nataf et al. 2012).

The final released data of Nataf et al. (2012) contain 9019 sightlines, which are almost entirely within the range $-10^\circ < l < 10^\circ$ and $2^\circ < |b| < 7^\circ$ and have a sky coverage of 90.25 deg^2 . The released catalogue contains the total number counts, mean distance modulus, geometrical variance, i.e. square of the dispersion in distance modulus, and full error matrix for each sightline.

These new data address two shortcomings in the study by Rattenbury et al. (2007). First, the total area in Rattenbury et al. (2007) covered only about 11 square degrees, spread over a longitude range of $-6.8^\circ \leq l \leq 10.6^\circ$ degrees. Furthermore, the field latitudes are clustered around (-4°) with only three fields at positive latitudes. Second, more seriously, Rattenbury et al. (2007) found that in order to match the luminosity of local and bulge red clump giants, an offset of 0.3 magnitude had to be artificially applied. The study by Nataf et al. (2012) addressed both issues. First, they resolve the “offset” difficulty by a much more careful consideration of extinction and a new calibration in the intrinsic luminosity of the red clump giants. Second, the number of fields is significantly more (9019), covering larger than 90 square degrees, nearly one order of magnitude larger than used in Rattenbury et al. (2007). As a result, the differences between different models become much more drastic and the constraints on the model parameters are tighter. The data also allow us to explore the residuals between data and model much more convincingly, which we will exploit in some detail (see §4).

3 METHODOLOGY

In this section, we first describe the parametric tri-axial models we use to model the luminosity density model of the Galactic bar, then we outline the Markov Chain Monte Carlo method we use to find the best-fit parameters for each model. The results of our procedure will be presented in the next section.

3.1 Parametric models

We first re-examine a few models of bar density distribution proposed by Dwek et al. (1995):

$$n_{G1} = N_0 \exp(-r^2/2), \quad (1)$$

$$n_{E2} = N_0 \exp(-r), \quad (2)$$

$$n_{E3} = N_0 K_0(r_s), \quad (3)$$

where K_0 is the modified Bessel function of the second kind and

$$r = \left[\left(\frac{x}{x_0} \right)^2 + \left(\frac{y}{y_0} \right)^2 + \left(\frac{z}{z_0} \right)^2 \right]^{1/2} \quad (4)$$

$$r_s = \left[\left[\left(\frac{x}{x_0} \right)^2 + \left(\frac{y}{y_0} \right)^2 \right]^2 + \left(\frac{z}{z_0} \right)^4 \right]^{1/4}. \quad (5)$$

These models are representative of a larger suite of parametric models studied by Rattenbury et al. (2007).

We set the origin of the coordinate system at the Galactic Centre (GC), with the x-y plane aligned with the mid-plane of the Galaxy and the z-direction parallel to the direction of the Galactic poles. The x-direction is defined to be co-linear with the semi-major axis of the bar. The functions are rotated by an angle α around the z-axis. N_0 is the number counts per kpc³, and the distance to the GC is another free parameter R_0 . Thus we have six parameters to model the bar structure.

The bar can also be rotated by a tilt angle β around the y axis, corresponding to the Sun's position away from the mid-plane of the Galaxy. Furthermore, there is tentative evidence that the centroid of the bar may be offset from the centre of mass of the Galaxy (as evidenced, for example, from the possible gamma-ray line emission offset from the centre, see Su & Finkbeiner 2012). We thus include two additional parameters in the modelling process, δl , δb , which allow for this possible centroid offset. We also fit the data using models in which all nine parameters are allowed to vary and compare the results with those found using the simpler six-parameter models.

With these models, we can calculate analytically the number counts, mean distance modulus and geometrical variance of each sightline as derived by Nataf et al. (2012).

The number counts in each sightline N_i is:

$$N_i = c_i \int_{s_{\min}}^{s_{\max}} n(s) s^2 ds, \quad (6)$$

where the constant c_i depends on the solid angle subtended around each sightline, and $s_{\min}$ and $s_{\max}$ are the minimum and maximum distances along the line of sight. We take $s_{\min} = 3$ kpc and $s_{\max} = 13$ kpc. We perform the integration over the range of $\sim R_0 \pm 5$ kpc as we do not expect the tri-axial bulge density structure to exceed these limits. The results are insensitive to the choice of integration limits since the bar density falls away rapidly and the contribution to the integral over the outer part of the bulge is negligible. Taking $\sim R_0 \pm 3$ kpc instead, the total number count changes only by 0.2% on average, and thus has the integration limit has little effect on our results.

The mean distance modulus of each sightline μ_i can be calculated as:

$$\mu_i = \int_{s_{\min}}^{s_{\max}} n(s) s^2 \mu ds / \int_{s_{\min}}^{s_{\max}} n(s) s^2 ds, \quad (7)$$

where the distance modulus is $\mu = 5 \log(s/\text{kpc}) + 10$.

The geometrical variance $\sigma_{\mu,i}$ can be written as,

$$\sigma_{\mu,i}^2 = \left[\int_{s_{\min}}^{s_{\max}} n(s) s^2 \mu^2 ds / \int_{s_{\min}}^{s_{\max}} n(s) s^2 ds \right] - \mu_i^2, \quad (8)$$

We can evaluate the χ^2 goodness-of-fit statistic using the number counts, mean distance modulus, geometrical variance, and their covariance matrix. The χ^2 of the bar model can be written as,

$$\chi^2 = \sum_i (\mathbf{X}_i^{\text{mod}} - \mathbf{X}_i^{\text{obs}})^T \mathbf{S}_i^{-1} (\mathbf{X}_i^{\text{mod}} - \mathbf{X}_i^{\text{obs}}). \quad (9)$$

$\mathbf{X}_i^{\text{mod}}$ comprises the number counts (N_i), mean distance modulus (μ_i) and geometrical variance ($\sigma_{\mu,i}^2$) for the i -th sightline for the model under consideration, while $\mathbf{X}_i^{\text{obs}}$ comprises the observed values of these variables for the same sightline. $\mathbf{S}_i$ is the covariance matrix of uncertainties. The data comprising $\mathbf{X}_i^{\text{mod}}$ are available from Nataf et al. (2012). The data include 9019 lines of sight in total, but not all are used, we exclude the ‘‘bad’’ sightlines which deviate far from the model (see below).

3.2 Markov Chain Monte Carlo search for best-fit models

For each of the parametric models, there are at least six parameters to constrain. We use Bayesian inference to provide a probabilistic distribution of tri-axial bar model parameters by combining prior information with the data. Information from the data and prior knowledge entirely determines the posterior probability distribution as follows:

$$\pi(x) \propto \mathcal{L}(\text{data}|x) \times \text{prior}(x). \quad (10)$$

We use Markov Chain Monte Carlo (MCMC) simulations to explore the parameter space and find the posterior distribution. MCMC can be implemented in a number of ways to explore the parameter space. In this work we use the standard random walk Metropolis-Hastings algorithm (Metropolis et al. 1953; Hastings 1970). At any point x_n in the chain we generate a new trial point x_{trial} by drawing parameter shifts from the symmetrical proposal distribution function $\phi(x - x_n)$. This algorithm is universal for any proposal distribution function. Here, we use a multi-dimensional Gaussian distribution in the vector of parameters. We accept a candidate state x_{n+1} with probability $\min\{1, \pi(x_{\text{trial}})/\pi(x_n)\}$, where $\pi(x)$ is the equilibrium distribution of the chain (Lewis & Bridle 2002).

For each model, the chain is started at the corresponding best-fit model values of parameter vector as given in Rattenbury et al. (2007). The model total star counts are normalised so that they reproduce the observed N_0 for each sightline. The MCMC

Model	Sightlines	R_0	x_0	y_0	z_0	α	β	δl	δb	χ_1^2	χ_2^2	χ_3^2	χ_4^2	$\chi^2/d.o.f$
G_1	7631	8.22	1.23	0.51	0.43	28.8				13801	15858	15349	2300	2.07
E_2	8118	8.17	0.68	0.28	0.26	31.7				12384	17710	11714	2691	1.83
E_3	8271	8.14	0.69	0.29	0.28	28.8				12316	18698	11302	937	1.74
G'_1	7749	8.27	1.24	0.52	0.42	29.1	-0.88	-0.31	-0.012	13995	16235	15513	2253	2.07
E'_2	8178	8.21	0.68	0.28	0.25	31.8	-4.73	-0.22	-0.023	12587	17902	11739	2168	1.81
E'_3	8196	8.20	0.81	0.33	0.30	32.6	-7.12	-0.21	-0.018	12825	18031	11314	2228	1.81

Table 1. The best fit values of six- and nine-parameter tri-axial galactic bar models.

simulation is then allowed to run, varying each of the six (or nine) parameters. Given the sampling strategy described above, we also use an adaptive step size to guarantee a sufficient sampling of the posterior, leading to convergence to the equilibrium posterior distribution in the chain after several thousand steps. We run 10^5 chains for each model to obtain the posterior distribution of the parameters, after neglecting the first half of the chain to allow the chain to reach its equilibrium state.

4 RESULTS

For the simple bar models, we have six parameters R_0, N_0, x_0, y_0, z_0 and α . We measure 2.94×10^6 RC stars in the OGLE-III data over a viewing area of 90.25 deg^2 in total. The total number of sightlines is 9019. For each model, we first run 10^5 Monte Carlo chains and get the best fit parameters for each model based on the original data set. We exclude sightlines that deviate more than 4σ , i.e. where $(\mathbf{X}_i^{\text{mod}} - \mathbf{X}_i^{\text{obs}})^T \mathbf{S}_i^{-1} (\mathbf{X}_i^{\text{mod}} - \mathbf{X}_i^{\text{obs}}) \geq 16$, from the model and then re-run to obtain the final results.

The results of best fit parameters are presented in Table 1. Since N_0 is not a true parameter and can be normalised by the total star counts, we include N_0 in the MCMC chain but do not give its value here. The number of sightlines for each model is also listed in Table 1. We separate χ^2 contributions from the number counts (χ_1^2), mean distance modulus (χ_2^2), geometrical variance (χ_3^2) respectively. χ_4^2 is the contribution from non-diagonal terms in the error matrix. Among these models, the E_2 and E_3 models have similar $\chi^2/d.o.f$ values and they are both much smaller than that of the G_1 model. This suggests that the E_2 and E_3 models best fit the data, while the G_1 model fits the data worst.

We present the marginalized posterior distribution of the parameters for the six-parameter G_1 model in Fig. 1. We can see that the probability distribution function (PDF) of each parameter is nearly Gaussian. We also show the degeneracy between each parameter in Fig. 1. The 1- and 2- σ regions (i.e., 68% and 95% confidence levels) are shown in inner and outer contour lines respectively. Fig. 5 is the same as Fig. 1, but for the E_2 model.

We compare the model prediction quantities, e.g. the surface density of stars Σ_{RC} , mean distance modulus μ , and geometrical variance σ_μ^2 , with that directly from observation for the models G_1 and E_2 . In the top left panel of Fig. 2, we plot Σ_{RC} for three stripes at $|b| = 2.5^\circ, 4.0^\circ, 5.5^\circ$ as observed in Nataf et al. (2012). Σ_{RC} is maximized near $l = 0^\circ$ for all latitude stripes. In the same figure, we also plot the G_1 model predictions of Σ_{RC} . At small longitudes, the model predictions are consistent with the observations. However, at large longitudes, especially for the $|b| = 5.5^\circ$ stripe, the model predictions are systematically lower than what is observed. In the top right panel of Fig. 2, we plot Σ_{RC} from both model predictions and observations at fixed longitudes of $l = 0.0^\circ, 5.0^\circ$ and 10.0° respectively. It shows the expected result that number counts increase monotonically with decreased separation from the plane for each longitude. However, it again shows that at large longitude, especially for the $l = 10.0^\circ$ stripe, the model predictions Σ_{RC} are systematically lower than the observed values.

The middle left panel of Fig. 2 shows the G_1 model (black symbols) and observed (red symbols) mean distance modulus μ against longitude. In the middle right panel, we plot μ for both model prediction and observation at fixed longitudes of $l = -5.0^\circ, 0^\circ$ and 5.0° . In the bottom panels of Fig. 6, we plot the modelled (black symbols) and observed (red symbols) geometrical variance σ_μ^2 vs. longitude (left) and latitude (right) respectively.

Notice that negative values of σ_μ^2 should not be a cause of concern. Nataf et al. (2012) estimated the dispersion due to geometric extension σ_μ^2 as:

$$\sigma_\mu^2 = \sigma^2 - \sigma_I^2 - \sigma_E^2 \quad (11)$$

where σ^2 is the measured brightness dispersion of the RC, $\sigma_I = 0.09$ is the estimated intrinsic magnitude dispersion of RC, and σ_E^2 is the dispersion due to differential extinction. Therefore, for sightlines where the measured brightness dispersion is small relative to the intrinsic dispersion and differential extinction, we can sometimes obtain negative variance in σ_μ^2 . We do not remove these sightlines with negative geometrical variance to avoid systematically biasing the final output to have large variance. However, in the bottom panels of Figs. 2 and 6 the predicted σ_μ^2 are well concentrated while the data have very wide variations. It could be that the models are wrong, or it could be that the data are wrong. To test for the latter, we take the intrinsic dispersion of the RC as a free parameter instead of $\sigma_I = 0.09$ as quoted in Nataf et al. (2012). Though very

Model E_2 with cut	Sightlines	R_0	x_0	y_0	z_0	α	χ^2_1	χ^2_2	χ^2_3	χ^2_4	$\chi^2/d.o.f$
$ l \leq 4^\circ$	5439	8.17	0.80	0.25	0.24	31.4	8127	11525	8166	1039	1.77
$ l \leq 4^\circ, b \leq 4^\circ$	4033	8.18	0.71	0.24	0.26	30.8	5432	9085	5472	237	1.67

Table 2. The best fit values of tri-axial galactic bar model after removing the outer part of the bulge regions.

different values of σ_I are suggested for different models; the models show little improvement in the $\chi^2/d.o.f$, and thus will not be discussed any further.

Comparing Fig. 6 with Fig. 2 we can see that a clear improvement over the G_1 model is that at large longitude, e.g. for the $|b| = 5.5^\circ$ stripe, the model surface density of stars Σ_{RC} is more consistent with the observed data.

In Fig. 3, we plot the mean distance to RC centroids measured by Nataf et al. (2012) and give the G_1 model predictions in the same figure for comparison. The distances are projected onto a face-on view of the central region of the Galaxy. The left panel of Fig. 3 shows data for sightlines with $2.5^\circ \leq |b| \leq 3.5^\circ$, while the right panel is for sightlines with $4.5^\circ \leq |b| \leq 5.5^\circ$. Fig. 7 is the same as Fig. 3, but for the E_2 model. The E_2 model is broadly consistent with the observations, but does not yield a very significant flattening at small positive longitudes as is seen in the data.

An interesting phenomenon, recently discovered, is that the RC of the bulge is observed to show two peaks in distance at large separations from the plane (Nataf et al. 2010; McWilliam & Zoccali 2010; Saito et al. 2011), a result that has been shown to be due to an X-shape structure by means of N-body models (Li & Shen 2012; Ness et al. 2012). The existence of the vertical X-shaped structure hints that the formation of the Milky Way bulge is shaped mainly by internal disk dynamical instabilities.

To verify the X-shaped fine structure, we calculate the excess fraction in star counts between the model and the observed data, $(N_{\text{obs}} - N_{\text{model}})/N_{\text{model}}$, for each sightline. We then project the excess fraction to produce a 2-D residual map, shown in the top panel of Fig. 4 for G_1 model. We can see a dim X-shaped structure extending from $l = 13^\circ$ to $l = -10^\circ$, with latitudes $|b| \leq 6^\circ$. After removing the bad sightlines, we still have 2.52×10^6 stars remaining. As we find a best-fit predicted RC population for the bulge of 2.35×10^6 , the excess RC population is 7.3%. In the bottom panel of Fig. 4, we also present the residual map of mean distance modulus. The regions have larger number counts show larger distances in observations than in models, but this trend does not hold on a pixel-by-pixel basis. Fig. 8 is the same as Fig. 4, but for the E_2 model. The predicted RC population for the bulge is 2.35×10^6 while the observed value is 2.52×10^6 with an excess of 5.3%. This value is not inconsistent with Li & Shen (2012), who found the excess light to be 7% in simulation. The agreement may be somewhat coincidental since their values are for an edge-on view. The existence of an X-shape structure suggests that at the outer regions of the bulge, there is some fine structure in the Galactic bar.

We also test the results of these models after introducing a centroid offset ($\delta l, \delta b$) and tilt angle (β). To avoid confusion, these models are labelled as G'_1, E'_2 and E'_3 . The results of the nine-parameter model fitting are also shown in Table 1. However, for the G'_1 and E'_2 model, the results of fitting a nine-parameter model show little improvement over the simpler six-parameter models. For the nine-parameter E'_3 model, a very slightly larger $\chi^2/d.o.f$ is obtained due to the more sightlines being included. This is perhaps not surprising since our sightlines do not cover the very central regions which presumably show the most prominent deviations from a non-offset model. Thus from now on $\beta, \delta l$ and δb will all be fixed at zero.

4.1 Disk contamination

There is evidence that there is some structural change in the Galactic bulge inward of $l \leq 4^\circ$ (Gerhard & Martinez-Valpuesta 2012). As mentioned above, the vertical X-shaped structure is assumed to be shaped mainly by internal disk dynamical instabilities and exists at larger longitudes. Both features are a source of disk contamination which is difficult for our model to reproduce, especially in regions remote from the centre of the bulge. From Figs. 2 and 6, we can see that at larger longitude, the number counts and geometrical variance in observation also systematically deviate from the model prediction.

To see the effects of possible disk contamination, we simply exclude data from the outer part of the bulge. We adopt two cut criteria. Firstly, we remove sightlines with $|l| > 4^\circ$. Secondly, we retain only the sightlines whose longitude and latitude are both smaller than 4° . The value of 4 degrees is chosen because in this region, there is a slope change in the particle surface density, see, e.g., Fig. 2 of Gerhard & Martinez-Valpuesta (2012).

We test the results of the E_2 model for the above two cut criteria. The results are given in Table 2. The model fit has an obviously smaller $\chi^2/d.o.f$ than the fit without cut. We again calculate the model-predicted RC population, and find the excess reduces to 3.3% for cutting the area of $|l| > 4^\circ$ and 2.6% for cutting $|l| > 4^\circ$ and $|b| > 4^\circ$. These are not surprise since we can see from Figs. 6 and 8 that the larger dispersion and excess both lie in the outlier of the bulge region. It suggests that we can get a much more realistic inner bar structure after cutting the wide longitude sightlines. The cuts on the data partly avoid the disk contamination and fine structure, which are presently very difficult to handle.

4.2 Comparison with results from OGLE-II

Rattenbury et al. (2007) used RCG stars from 44 bulge fields from the OGLE-II microlensing collaboration database to constrain analytical tri-axial models for the Galactic bar. They found the bar major axis is oriented at an angle of 24° to 27° to the Sun-Galactic centre line of sight. They also found the ratio of semi-major and semi-minor bar axis scale lengths in the Galactic plane x_0 , y_0 , and vertical bar scale length z_0 , to be $x_0 : y_0 : z_0 = 1.00 : 0.35 : 0.26$.

As mentioned in the introduction, the data from OGLE-III as analyzed by Nataf et al. (2012) offer many more sightlines and better calibration in the RCG population. Both are helpful to refine the constraints on the bar parameters. We remove the area in the outer part of the bulge region and keep the fields with $-4^\circ \leq l \leq 4^\circ$ and $-4^\circ \leq b \leq 4^\circ$ to partly remove the disk contamination and concentrate on the inner region of the bar. When using all the data, we found an excess, consistent with the so-called the X-shape structure in the outer part of the bulge region. The excess is $\sim 7\%$, in agreement with Li & Shen (2012), but which complicated the fitting. When we remove data from the outer part of the bulge, the effect of the fine structure is reduced and we can fit a smoother inner bar structure. We obtain a bar angle oriented at 29° to 32° , and the axis ratios are around $x_0 : y_0 : z_0 = 1.00 : 0.41 : 0.38$. This suggests a bulge that is not as tri-axial as the bar structure found by Rattenbury et al. (2007); it is also vertically more extended than the working model of Gerhard (2002) with $x_0 : y_0 : z_0 = 1.00 : 0.40 : 0.30$.

5 DISCUSSION

We use the RCG population in the OGLE-III database to fit a tri-axial Galactic bar model and to derive much tighter constraints on the bar parameters. When using all the available data, we found there is evidence for an X-shaped structure in the outer part of the bulge region, with observable properties consistent with those of Li & Shen (2012). This fine structure complicated the fitting of our analytic models. We removed data from the area in the outer part of the bulge region and retained the fields with $-4^\circ \leq l \leq 4^\circ$ and $-4^\circ \leq b \leq 4^\circ$ to partly remove disk contamination and to concentrate on the inner region of the bar. This reduced the effect of the fine structure and allowed a fit to the resulting smoother inner bar structure. We obtain a bar angle between 29° to 32° with an axis ratio around $x_0 : y_0 : z_0 = 1.00 : 0.41 : 0.38$. This bar is close to being prolate, and not as triaxial as the bar structure found by Rattenbury et al. (2007); it appears vertically thicker than the working model of Gerhard (2002) who gives $x_0 : y_0 : z_0 = 1.00 : 0.40 : 0.30$. The bar is more prolate than that found by Dwek et al. (1995) and has a larger bar angle. The model derived here can effectively reproduce the number density, distance modulus over the sky probed by the OGLE-III data (Nataf et al. 2012).

Future microlensing observations will provide us with additional constraints on the structure of the Galactic bar. Based on the improved and tightly constrained bar model, it will be interesting to re-calculate the optical depths and time-scale distribution for microlensing events, and see whether they are consistent with the data.

The X-shaped structure appears in both simulations and in our data. Its existence in the Milky way implies that the Galactic bulge is shaped mainly by internal disk dynamical instabilities instead of mergers (e.g. Li & Shen 2012). How to model subtle structures such as the X-shaped excess remains a challenging task. One possible way forward is to expand the density distribution into orthogonal basis functions; a similar approach is taken in weak lensing to model complex point spread functions (Andrae et al. 2011); we plan to explore this in the near term.

ACKNOWLEDGMENTS

We acknowledge the National Astronomical Observatories (NAOC) and Chinese Academy of Sciences (CAS) for financial support (SM). CL acknowledges the support of the NSFC grant(11203028) and the Young Researcher Grant of NAOC. This paper is also made possible through the support of a grant from John Templeton Foundation. Andrew Gould acknowledges the NSF grant (AST-1103471) and is also partly supported by a senior visiting professorship of CAS. We thank Drs. Lia Athanassoula, Ortwin Gerhard and Guoliang Li for helpful discussions. CL also thanks Dr. Junqiang Ge for his help.

REFERENCES

- Andrae, R., Melchior, P., & Jahnke, K. 2011, MNRAS, 417, 2465
- Athanassoula, E. 2012, arXiv:1211.6752
- Babusiaux, C., Gilmore, G., 2005, MNRAS, 358, 1309
- Binney, J., Gerhard, O.E., Stark, A.A., et al., 1991, MNRAS, 252, 210
- Blitz, L., Spergel, D.N., 1991, ApJ, 379, 631
- Dwek, E., Arendt, R.G., Hauser, M.G., et al., 1995, ApJ, 445, 716
- Freeman, K., Ness, M., Wylie-de-Boer, E., et al. 2013, MNRAS, 428, 3660

- Gerhard, O., 2002, in G.S. Da Costa, H. Jerjen, eds., ASP Conf. Ser. 273: The Dynamics, Structure & History of Galaxies: A Workshop in Honour of Professor Ken Freeman, 73
- Gerhard, O., & Martinez-Valpuesta, I. 2012, ApJ, 744, L8
- Gonzalez, O. A., Rejkuba, M., Minniti, D., et al. 2011, A&A, 534, L14
- Hastings, W.K. 1970, Biometrika 57,97
- Howard, C. D., Rich, R. M., Reitzel, D. B., et al. 2008, ApJ, 688, 1060
- Lewis, A., & Bridle, S. 2002, Phys. Rev. D, 66, 103511
- Li, Z.-Y., & Shen, J. 2012, ApJ, 757, L7
- Long, R. J., Mao, S., Shen, J., & Wang, Y. 2013, MNRAS, 428, 3478
- McWilliam, A., & Zoccali, M. 2010, ApJ, 724, 1491
- Ness, M., Freeman, K., Athanassoula, E., et al. 2012, ApJ, 756, 22
- Saito, R. K., Zoccali, M., McWilliam, A., et al. 2011, AJ, 142, 76
- Metropolis, N.; Rosenbluth, A.W.; Rosenbluth, M.N.; Teller, A.H.; Teller, E. 1953. Journal of Chemical Physics 21, 1087
- Nakada, Y., Onaka, T., Yamamura, I., et al., 1991, Nature, 353, 140
- Nataf, D. M., Udalski, A., Gould, A., Fouqué, P., & Stanek, K. Z. 2010, ApJ, 721, L28
- Nataf, D. M., Gould, A., Fouqué, P., et al. 2012, arXiv:1208.1263
- Nidever, D. L., Zasowski, G., Majewski, S. R., et al. 2012, ApJ, 755, L25
- Nishiyama, S., Nagata, T., Baba, D., et al., 2005, ApJ, 621, L105
- Rattenbury, N. J., Mao, S., Sumi, T., & Smith, M. C. 2007, MNRAS, 378, 1064
- Sellwood J. A., Wilkinson A., 1993, RPPh, 56, 173
- Stanek, K.Z., Mateo, M., Udalski, A., et al., 1994, ApJ, 429, L73
- Stanek, K.Z., Udalski, A., Szymanski, M., et al., 1997, ApJ, 477, 163
- Su, M., & Finkbeiner, D. P. 2012, arXiv:1206.1616
- Terndrup, D. M., Popowski, P., Gould, A., Rich, R. M., & Sadler, E. M. 1998, AJ, 115, 1476
- Udalski, A., Szymanski, M., Stanek, K.Z., et al., 1994, Acta Astronomica, 44, 165
- Wang, Y., Zhao, H., Mao, S., & Rich, R. M. 2012, MNRAS, 427, 1429
- Weiland, J.L., Arendt, R.G., Berriman, G.B., et al., 1994, ApJ, 425, L81




Figure 1. The marginalised probability density distributions (PDFs) and contours in the plane of R_0 , x_0 , y_0 , z_0 , α for the six-parameter G_1 model. All the lengths are in units of kpc and the angle α is measured in degrees. The inner and outer contours are respectively for the $1\text{-}\sigma$ and $2\text{-}\sigma$ confidence level regions (for two parameters simultaneously).




Figure 2. The predicted surface density of stars Σ_{RC} , mean modulus μ , and geometrical variance σ_μ^2 , vs. those directly from observation G_1 model. Top panels: the surface number density Σ_{RC} vs. longitude (left) and latitude (right). Black symbols denote model values while colour symbols denote observational data. Middle panels: mean distance modulus μ vs. longitude (left) and latitude (right). In the middle and bottom panels, black symbols denote model values and red symbols observational data. Bottom panels: geometrical variance σ_μ^2 vs. longitude (left) and latitude (right).




Figure 3. The face-on view projection of the distances to RC centroids for the G1 model. The red points show the projected location of each measured RC centroid, and black points show the model prediction. The left panel is for sightlines satisfying $2.5^\circ \leq |b| \leq 3.5^\circ$, while the right panel is for $4.5^\circ \leq |b| \leq 5.5^\circ$ sightlines.




Figure 4. The residual map of number counts (top) and mean distance modulus (bottom) for the six-parameter G_1 model. The residual of number counts is defined as the fractional difference in star counts between the model and the observed data, $(N_{\text{obs}} - N_{\text{model}})/N_{\text{model}}$. The total excess fraction of number counts is 7.3%. The residual in the mean distance is defined as $\mu_{\text{obs}} - \mu_{\text{model}} = 5 \log D_{\text{obs}}/D_{\text{model}}$, where D_{obs} and D_{model} are the the observed and predicted distances.




Figure 5. The same as Fig. 2, but for the six-parameter E_2 model.




Figure 6. The same as Fig. 2, but for the six-parameter E_2 model.




Figure 7. The same as Fig. 3, but for the six-parameter E_2 model.




Figure 8. The same as Fig. 4, but for the six-parameter E_2 model. The total excess fraction of number counts is 5.3%.